arXiv:1201.1845v4 [cond-mat.other] 10 Nov 2012

Microstructure of He II in the presence of boundaries

M. D. Tomchenko*
Bogolyubov Institute for Theoretical Physics, 14-b Metrolohichna Street, Kiev 03680, Ukraine

We show that, in spite of the commonly accepted expectations, the boundaries affect strongly the bulk microstructure of He II. We determine the structure of exact wave functions of the ground (Ψ_0) and excited ($\Psi_{\mathbf{k}}$) states of He II under zero boundary conditions. The explicit account for boundaries leads to a significant change of the equations for Ψ_0 and $\Psi_{\mathbf{k}}$ and the theoretical values of the energy of quasiparticles $E(k)$ and the energy of the ground state E_0, as compared with the solutions for periodic boundary conditions. In the limit of a weak interaction, we obtain the formula $E(k) \approx \sqrt{\left(\frac{\hbar^2k^2}{2m}\right)^2 + \frac{n\nu_3(k)}{2^d}\frac{\hbar^2k^2}{m}}$ (d is the space dimensionality), which differs from the well-known Bogolyubov formula by the factor 2^{-d}. The effect of boundaries is topological and is related to the different collections of eigenmodes for closed and open system.

PACS numbers: 67.10.-j, 67.25.dj, 67.25.dt

I. INRODUCTION

Starting from the pioneer works by L. Landau [1], N. Bogolyubov [2, 3], and R. Feynman [4, 5], the microscopic theory of superfluid He^4 (He II) is developed for about seven decades. It explains sufficiently well the basic properties of He II (see reviews [6–8]) and is considered to be established. At the same time, some questions remain open. In particular, the nature of the λ-transition, number of higher condensates, properties and role of microscopic vortex rings are not clear. Moreover, the relationship of superfluidity to the presence of a condensate of atoms is not completely clarified.

In addition, the construction of a microscopic model uses always the periodic boundary conditions (BCs) with the subsequent transition to the thermodynamic limit ($N, V \to \infty$, $N/V = const$) instead of realistic BCs (a finite system with the surface). Certain difficulties in the construction of a perturbation theory at realistic BCs were indicated in the articles by A. Bijl [9] and M. Girardeau [10] and the monograph by K. Huang [11]. This is due to the fact that the bare wave function (WF) is naturally sought in the form of a solution for the ideal gas, as a product of sines $\sin(\pi x_j/L_x)$ (in the one-dimensional case). But such sines set a very inhomogeneous "arc-like" density of a medium, whereas the density of real helium is homogeneous. In view of this difficulty, the realistic BCs were changed by the periodic ones in order to simplify the calculations.

To our knowledge, the solutions for the WFs of interacting Bose particles with the explicit account for boundaries were obtained only for a one-dimensional system of particles with point interaction [12, 13]. We will find the solutions for particles with any nonpoint interatomic potential possessing the Fourier-transform and for any dimensionality.

It is clear intuitively that the direct influence of boundaries on a bulk Bose fluid (e.g., He II) should be small. But He II is a strongly nonlinear and highly correlated system with a condensate. Since the condensate corresponds to a delicate tuning of the whole system, even a very weak effect of boundaries can change it. We will show that the boundaries strongly affect a quantum fluid: they affect the spectrum of eigenmodes of the ground and excited states. As a result, the calculated values of the ground state (GS) energy, energy of quasiparticles, and a structure of a condensate [14] become essentially different from those for periodic BCs.

II. GROUND STATE OF HE II: EQUATIONS

Several methods of determination of the WF Ψ_0 of the GS of helium-II [3, 15–26] (see also reviews [7, 8]) are available. They used periodic BCs, which cannot be obviously realized in the nature. Our task is to find Ψ_0 under BCs close to those realized in the nature and for realistic interatomic potentials. The influence of boundaries on the structure of the WF can be nontrivial. Therefore, it is expedient to use one of the exact analytic methods for the determination of Ψ_0 [15–17, 20], since the numerical methods may not catch the effect. We will operate within the method of collective variables [17, 27] used by us [25, 28].

We transit to the collective variables:

$$\rho_{\mathbf{k}} = \frac{1}{\sqrt{N}} \sum_{j=1}^{N} e^{-i\mathbf{k}\mathbf{r}_j} \quad (\mathbf{k} \neq 0), \tag{1}$$

where N is the total number of atoms, and $\mathbf{r}_j$ are the coordinates of atoms. For convenience, we introduce the following notation. The sums over the wave vectors

$$\mathbf{k} = 2\pi \left(\frac{j_x}{L_x}, \frac{j_y}{L_y}, \frac{j_z}{L_z}\right) \tag{2}$$

(j_x, j_y, j_z are integers, and L_x, L_y, L_z are the sizes of the

---

*Electronic address: mtomchenko@bitp.kiev.ua

system) are denoted by the index (2π); for example, $\sum\limits_{\mathbf{k}}^{(2\pi)}$. And the sums over the wave vectors

$$\mathbf{k} = \pi\left(\frac{j_x}{L_x}, \frac{j_y}{L_y}, \frac{j_z}{L_z}\right) \tag{3}$$

are denoted by the index (π). For example, $\sum\limits_{\mathbf{k},\mathbf{q}\neq 0}^{(\pi)}$ means the sum over $\mathbf{k}$ and $\mathbf{q}$, which are nonzero and run series (3). If the row with (π) or (2π) contains some designation (e.g., $\mathbf{k}+\mathbf{q}\neq 0$), it has no relation to (π) or (2π).

At the periodic BCs, the solution for Ψ_0 is obtained by I. Yukhnovskii and I. Vakarchuk [17]:

$$\Psi_0 = Ae^{S_b}, \tag{4}$$

$$\begin{aligned} S_b &= \sum_{\mathbf{k}_1\neq 0}^{(2\pi)} \frac{a_2(\mathbf{k}_1)}{2!}\rho_{\mathbf{k}_1}\rho_{-\mathbf{k}_1} + \\ &+ \sum_{\mathbf{k}_1,\mathbf{k}_2\neq 0}^{(2\pi)\mathbf{k}_1+\mathbf{k}_2\neq 0} \frac{a_3(\mathbf{k}_1,\mathbf{k}_2)}{3!\sqrt{N}}\rho_{\mathbf{k}_1}\rho_{\mathbf{k}_2}\rho_{-\mathbf{k}_1-\mathbf{k}_2} + \\ &+ \sum_{\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3\neq 0}^{(2\pi)\mathbf{k}_1+\mathbf{k}_2+\mathbf{k}_3\neq 0} \frac{a_4(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3)}{4!N}\rho_{\mathbf{k}_1}\rho_{\mathbf{k}_2}\rho_{\mathbf{k}_3}\rho_{-\mathbf{k}_1-\mathbf{k}_2-\mathbf{k}_3} \\ &+ \ldots, \end{aligned} \tag{5}$$

where the functions $a_j(\mathbf{k}_1,\ldots,\mathbf{k}_{j-1})$ satisfy the chain of equations

$$E_0^b = \frac{N-1}{2N}n\nu_3(0) - \sum_{\mathbf{k}\neq 0}^{(2\pi)}\frac{n\nu_3(k)}{2N} - \sum_{\mathbf{k}\neq 0}^{(2\pi)}\frac{\hbar^2k^2}{2mN}a_2(\mathbf{k}), \tag{6}$$

$$\begin{aligned} &\frac{n\nu_3(k)m}{\hbar^2} + a_2(\mathbf{k})k^2 - a_2^2(\mathbf{k})k^2 = \\ &= \sum_{\mathbf{q}\neq 0}^{(2\pi)}\frac{a_3(\mathbf{k},\mathbf{q})}{N}(q^2+\mathbf{k}\mathbf{q}) + \sum_{\mathbf{q}\neq 0}^{(2\pi)} q^2\frac{a_4(\mathbf{q},-\mathbf{q},\mathbf{k})}{2N}, \end{aligned} \tag{7}$$

$$a_3(\mathbf{k},\mathbf{q}) \approx -2\frac{R(\mathbf{k},\mathbf{q})}{\epsilon_0(k)+\epsilon_0(q)+\epsilon_0(\mathbf{k}+\mathbf{q})}, \tag{8}$$

$$\begin{aligned} R(\mathbf{k},\mathbf{q}) &= \mathbf{k}\mathbf{q}a_2(\mathbf{k})a_2(\mathbf{q}) - \mathbf{k}(\mathbf{k}+\mathbf{q})a_2(\mathbf{k})a_2(\mathbf{k}+\mathbf{q}) - \\ &- \mathbf{q}(\mathbf{k}+\mathbf{q})a_2(\mathbf{q})a_2(\mathbf{k}+\mathbf{q}), \end{aligned} \tag{9}$$

$$\epsilon_0(\mathbf{k}) = k^2(1-2a_2(\mathbf{k})), \tag{10}$$

where $n = N/V$, V is the volume of the system, E_0^b is the "bulk" energy of the GS per atom, and

$$\nu_3(\mathbf{k}) \equiv \nu_3(k) = \int U_3(r)e^{-i\mathbf{k}\mathbf{r}}d\mathbf{r} \tag{11}$$

is the Fourier-transform of the interaction potential $U_3(r)$ of two He^4 atoms. The equation for a_3 is written in the zero approximation not involving the corrections a_4 and a_5, and the equations for higher corrections $a_{j\geq 4}(\mathbf{k}_1,\ldots,\mathbf{k}_{j-1})$ are not presented.

Consider He II in a rectangular vessel $L_x \times L_y \times L_z$ in size. In order to solve the problem exactly, the solution should be sought inside and outside of the vessel, and the continuity of the WF and all its derivatives on the boundary can be used as the BCs. However, it is too complicated. To simplify, let us consider impermeable walls. Then the WF is nonzero inside the vessel, and outside of it and on the boundary is equal to zero. The real walls are not impermeable, but they create a high potential barrier with $U_s \gtrsim 1000\,K$ (see Eq. (14)). At such a barrier, the solution for a single particle in the box differs from the solution under an infinitely high (impermeable) barrier arbitrarily slightly.

The Hamiltonian of liquid helium bounded by the walls has the form

$$\hat{H} = -\frac{\hbar^2}{2m}\sum_j \triangle_j + \frac{1}{2}\sum_{ij}^{i\neq j} U_3(|\mathbf{r}_i-\mathbf{r}_j|) + U_2 + U_{em}, \tag{12}$$

$$\begin{aligned} U_2 &= \sum_j [U_w(x_j) + U_w(L_x-x_j) + U_w(y_j) + \\ &+ U_w(L_y-y_j) + U_w(z_j) + U_w(L_z-z_j)]. \end{aligned} \tag{13}$$

Here, U_2 is the interaction potential of helium atoms with all walls. The potentials of the walls are similar by form [29] to the interaction potential of two helium atoms. The free surface of liquid helium creates an energy barrier also. Therefore, we consider it as a wall. Below, we assume all barriers identical with simple potential

$$U_w(x) = \left[\begin{array}{ll} U_s > 0 & x \leq 0, \\ 0 & x > 0. \end{array}\right. \tag{14}$$

Below, we take U_s finite, and the transition $U_s \to \infty$ will be made in final formulas.

Works [17, 27] used the non-Hermitian Bogolyubov–Zubarev Hamiltonian. Since the absence of the Hermitian property induces the questions [30], we will start from the ordinary Hermitian Hamiltonian (12).

As is known, a helium atom disturbs the distribution of charges in a wall. If a wall is metallic, the perturbation is equivalent to an exact mirror "image" of the atom in the wall. In the image, the nucleus and "electrons" have, respectively, negative and positive charges. If the wall is made of a dielectric, the charges of the image are diminished. The interaction with images is taken into account in U_{em}. However, the consideration of the images strongly complicates the calculations. Therefore, we set $U_{em} = 0$ in what follows.

We expand the potential in a Fourier series:

$$U_3(|\mathbf{r}_1-\mathbf{r}_2|) = \frac{1}{2^dV}\sum_{\mathbf{k}}^{(\pi)}\nu_3(\mathbf{k})e^{i\mathbf{k}(\mathbf{r}_1-\mathbf{r}_2)}, \tag{15}$$

where $d = 3$ is the space dimensionality. We seek the WF of helium atoms inside the vessel, since the WF is zero outside of it. In this case, all functions in the Schrödinger equation should be expanded in a Fourier series with regard for the fact that the coordinates $\mathbf{r}_j$ of atoms are defined only in the interior of the vessel. As is known, such an expansion reproduces exactly the values of functions for all $\mathbf{r}_j$ inside of the vessel. In the potential $U_3(|\mathbf{r}_i - \mathbf{r}_j|)$, the coordinates x_i and x_j take values from the interval $[0, L_x]$, whereas the values of $x_i - x_j$ belong to the interval $[-L_x, L_x]$. Analogously, we consider the y- and z-components. By the rules of Fourier analysis, the function $U_3(|\mathbf{r}_i - \mathbf{r}_j|)$ must be expanded in this case in series (15): with $\mathbf{k}$ (3) and the volume equal to $2L_x \times 2L_y \times 2L_z = 2^d V$.

For the total interatomic potential, we obtain

$$\begin{aligned} \frac{1}{2}\sum_{i\neq j} U_3(|\mathbf{r}_i - \mathbf{r}_j|) &= \sum_{\mathbf{k}\neq 0}^{(\pi)} \frac{n\nu_3(k)}{2^{d+1}}\rho_{\mathbf{k}}\rho_{-\mathbf{k}} + \\ +\ (N-1)\frac{n\nu_3(0)}{2^{d+1}} &- \sum_{\mathbf{k}\neq 0}^{(\pi)} \frac{n\nu_3(k)}{2^{d+1}}. \end{aligned} \tag{16}$$

Under periodic BCs, $x_i - x_j$ take values from the interval $[0, L_x]$. Therefore, we need to replace $2^d V$ in (15) by V and to quantize $\mathbf{k}$ by law (2). Then we have [3]

$$\begin{aligned} \frac{1}{2}\sum_{i\neq j} U_3(|\mathbf{r}_i - \mathbf{r}_j|) &= \sum_{\mathbf{k}\neq 0}^{(2\pi)} \frac{n\nu_3(k)}{2}\rho_{\mathbf{k}}\rho_{-\mathbf{k}} + \\ +\ (N-1)\frac{n\nu_3(0)}{2} &- \sum_{\mathbf{k}\neq 0}^{(2\pi)} \frac{n\nu_3(k)}{2}. \end{aligned} \tag{17}$$

This is a key point. Since Eqs. (5)–(8) are "generated" by the total potential, we may assert at once that the explicit account for boundaries will lead to the replacement (in these equations) of $\nu_3(k)$ by $\nu_3(k)/2^d$ and the summation law (2) by (3).

Let us determine the WF Ψ_0 for the system of Bose particles in a vessel. Ψ_0 satisfies the Schrödinger equation

$$\hat{H}\Psi_0 = NE_0\Psi_0 \tag{18}$$

with Hamiltonian (12)–(16), where we neglect U_{em}.

The main question is in which form we should seek Ψ_0. The solution of (4), (5) for Ψ_0 with periodic BCs is usually obtained with regard for the requirement of the translation invariance and the functional independence [17] of the following collections of variables $\rho_{\mathbf{k}}$:

$$\rho_{-\mathbf{k}_1}, \quad \rho_{-\mathbf{k}_1}\rho_{-\mathbf{k}_2}, \quad \rho_{-\mathbf{k}_1}\rho_{-\mathbf{k}_2}\rho_{-\mathbf{k}_3}, \quad \ldots \tag{19}$$

But the boundaries break the translation invariance. However, the solution for (4), (5) can be obtained without regard for the translation invariance, by substituting the bare function $\Psi_0 = const$, describing the free bosons, in the Schrödinger equation. The equation itself "sets" the form of corrections, which appear in the exponent. For the problem with boundaries, we will seek Ψ_0 analogously, by substituting the solution for free particles in the Schrödinger equation.

The WFs of the system of N free Bose particles located in a rectangular box with potential (13), (14) are known. If the states of all particles are identical, the WF is equal to one of the functions [31]

$$\Psi_{\mathbf{l}}^f = \prod_{j=1}^{N}\{\sin(k_{l_x}x_j + \delta_{l_x})\sin(k_{l_y}y_j + \delta_{l_y})\sin(k_{l_z}z_j + \delta_{l_z})\}, \tag{20}$$

where $l_x, l_y, l_z = 1, 2, 3, \ldots$, and k_{l_x} and δ_{l_x} satisfy the equations

$$\sin\delta_{l_x} = \frac{\gamma_x k_{l_x} L_x}{2}, \quad \sin(k_{l_x}L_x + \delta_{l_x}) = \pm\frac{\gamma_x k_{l_x} L_x}{2}, \tag{21}$$

which yields

$$k_{l_x}L_x = \pi l_x - 2\delta_{l_x}, \quad \delta_{l_x} = \arcsin(\gamma_x k_{l_x} L_x/2). \tag{22}$$

Here, we denote $\gamma_x = \frac{\hbar}{L_x}\sqrt{\frac{2}{mU_s}}$, and the values of arcsin are taken between 0 and $\pi/2$. For the realistic systems, $\gamma_x \ll 1$. For example, for He^4 atoms at $U_s = 100\,K$ and $L_x = 10\,\mathrm{cm}$ we have $\gamma_x \approx 4.92 \times 10^{-10}$. At $l_x \ll 1/\pi\gamma_x$, relation (22) yields

$$k_{l_x} \approx l_x k_{1x} = l_x(1 - \gamma_x)\pi/L_x, \tag{23}$$

$$\delta_{l_x} \approx l_x\delta_{1x} = l_x\pi\gamma_x/2. \tag{24}$$

The maximally possible values of δ_{l_x} and k_{l_x} are determined from the condition $\sin\delta_{l_x} = 1$:

$$\delta_{l_x}^{max} = \pi/2, \quad k_{l_x}^{max} = k_{1x}/\delta_{1x}. \tag{25}$$

The WF of the ground state of a single atom inside of the box,

$$\Psi_{0,1}^f(\mathbf{r}) = \sin(k_{1x}x + \delta_{1x})\sin(k_{1y}y + \delta_{1y})\sin(k_{1z}z + \delta_{1z}),$$

is positive and nonzero. Outside of the box, the WF decays exponentially.

With regard for the above consideration, the WF of the ground state of N interacting Bose particles located in the box can be sought in the form

$$\Psi_0 = A_1\Psi_{\mathbf{0}}^f e^{S_w^{(1)} + \tilde{S}_b}, \tag{26}$$

where $\Psi_{\mathbf{0}}^f$ is the WF of the ground state of N free Bose particles in the box (function (20) with $l_x = l_y = l_z = 1$), $e^{\tilde{S}_b}$ is a WF of the form (4), (5) with the sums $\sum^{(\pi)}$ instead of $\sum^{(2\pi)}$ (this is seen from expansion (16)), and $e^{S_w^{(1)}}$ describes the cross terms (from $\Psi_{\mathbf{0}}^f$ and $e^{\tilde{S}_b}$) arising at the substitution of $\Psi_{\mathbf{0}}^f e^{\tilde{S}_b}$ in the Schrödinger equation. By the theorem on nodes, if the GS is nondegenerate, then Ψ_0 is nonzero in all points in the box. Therefore, it is convenient to lift the function $S_w^{(1)}$ to the exponent. The factor $e^{\tilde{S}_b}$ appears due to the interatomic interaction, like that in (4). We note that $\Psi_{\mathbf{0}}^f$ will ensure the fulfillment of BCs and the transition to the WF of free particles at the switch-off of the interaction. This gives the structure of (26).

Notice that the structure (26) was proposed previously by S. Yushchenko [32].

The function Ψ_0 (26) must satisfy the Schrödinger equation (18) and be zero at the boundaries. The last property is ensured at $U_s \to \infty$ by the factor $\Psi_{\mathbf{0}}^f$ which satisfies the equations

$$-\frac{\hbar^2}{2m}\sum_j \triangle_j \Psi_{\mathbf{0}}^f + U_2\Psi_{\mathbf{0}}^f = NE_0^f\Psi_{\mathbf{0}}^f, \tag{27}$$

$$E_0^f = \frac{\hbar^2 k_1^2}{2m}, \quad k_1^2 = k_{1x}^2 + k_{1y}^2 + k_{1z}^2. \tag{28}$$

The function $\tilde{S}_b$ satisfies the equation

$$-\frac{\hbar^2}{2m}\sum_j \triangle_j e^{\tilde{S}_b} + \frac{1}{2}\sum_{i\neq j} U_3(|\mathbf{r}_i - \mathbf{r}_j|)e^{\tilde{S}_b} = N\tilde{E}_0^b e^{\tilde{S}_b}. \tag{29}$$

Then Eq. (18) is reduced to the equation for $S_w^{(1)}$:

$$\begin{aligned} &- \frac{\hbar^2}{2m}\sum_j \Bigg[\triangle_j S_w^{(1)} + (\nabla_j S_w^{(1)})^2 + 2\nabla_j S_w^{(1)}\frac{\nabla_j \Psi_{\mathbf{0}}^f}{\Psi_{\mathbf{0}}^f} + \\ &+ \ 2\nabla_j \tilde{S}_b\left(\nabla_j S_w^{(1)} + \frac{\nabla_j \Psi_{\mathbf{0}}^f}{\Psi_{\mathbf{0}}^f}\right)\Bigg] = \\ &= \ N(E_0 - \tilde{E}_0^b - E_0^f). \end{aligned} \tag{30}$$

We note that

$$\begin{aligned} \frac{\nabla_j \Psi_{\mathbf{0}}^f}{\Psi_{\mathbf{0}}^f} &= \mathbf{i}_x k_{1x}\cot{(k_{1x}x_j + \delta_{1x})} + \\ &+ \quad \mathbf{i}_y k_{1y}\cot{(k_{1y}y_j + \delta_{1y})} + \mathbf{i}_z k_{1z}\cot{(k_{1z}z_j + \delta_{1z})}. \end{aligned} \tag{31}$$

By seeking Ψ_0 inside of the vessel, we set $\cot{(k_{1x}x + \delta_{1x})}$ on the interval $[0, L_x]$. In order to apply the method of collective variables, we expand the cotangent in the Fourier series:

$$\cot{(k_{1x}x + \delta_{1x})} = \sum_{j_x}^{(2\pi)} C_{1x}(j_x)e^{i2\pi j_x x/L_x}, \tag{32}$$

where j_x runs all integers. We have

$$\begin{aligned} C_{1x}(j_x) = -C_{1x}(-j_x) &= \int\limits_0^{L_x}\frac{dx}{L_x}\cot{(k_{1x}x + \delta_{1x})}e^{-iq_x x} \\ = \ -2i\int\limits_0^{1/2} dx \sin{(2\pi j_x x)} &\cot{(\pi x - 2\delta_{1x}x + \delta_{1x})}. \end{aligned} \tag{33}$$

Whence $C_{1x}(0) = 0$, and the function $C_{1x}(j_x)$ is approximated by the formula

$$C_{1x}(j_x > 0) \approx -i + \frac{i4j_x\delta_{1x}}{\pi}\ln\left(1 + \frac{\pi}{4j_x\delta_{1x}}\right), \tag{34}$$

which is valid at $j_x \lesssim j_m = 1/\delta_{1x}$ with an error of $\leq 3\%$.

We note that $\cot{(k_{1x}x + \delta_{1x})}$ satisfies the requirements to functions expanded in the Fourier series. At $\delta_{1x} = 0$, we obtain $\cot{(\pi x/L_x)}$, which does not satisfy these requirements, since the integral of the modulus of this function over the interval $[0, L_x]$ is equal to infinity. In view of this difficulty, we preserve all δ to be nonzero in calculations and then set $\delta \to 0$ in final formulas.

Using the difference of the functional structure of the collections $\rho_{\mathbf{k}}$ (19), we determine the structure of the solution for $S_w^{(1)}$ from (30)–(32) and (5):

$$\begin{aligned} S_w^{(1)} &= \sum_{\mathbf{q}\neq 0}^{(2\pi)} S_1^{(1)}(\mathbf{q})\rho_{-\mathbf{q}} + \sum_{\mathbf{q},\mathbf{q}_1\neq 0}^{\mathbf{q}+\mathbf{q}_1\neq 0}\frac{S_2^{(1)}(\mathbf{q},\mathbf{q}_1)}{\sqrt{N}}\rho_{\mathbf{q}_1}\rho_{-\mathbf{q}_1-\mathbf{q}} \\ &+ \sum_{\mathbf{q},\mathbf{q}_1,\mathbf{q}_2\neq 0}^{\mathbf{q}+\mathbf{q}_1+\mathbf{q}_2\neq 0}\frac{S_3^{(1)}(\mathbf{q},\mathbf{q}_1,\mathbf{q}_2)}{N}\rho_{\mathbf{q}_1}\rho_{\mathbf{q}_2}\rho_{-\mathbf{q}_1-\mathbf{q}_2-\mathbf{q}} + \ldots \end{aligned} \tag{35}$$

where $\mathbf{q}$ and $\mathbf{q}_j$ run values (2) and (3), respectively (the first fact follows from (32), and the second does from (15) and (7)). The functions $S_j^{(1)}$ satisfy the symmetry relations

$$S_2^{(1)}(\mathbf{q},\mathbf{q}_1) = S_2^{(1)}(\mathbf{q},-\mathbf{q}-\mathbf{q}_1), \tag{36}$$

$$\begin{aligned} S_3^{(1)}(\mathbf{q},\mathbf{q}_1,\mathbf{q}_2) &= S_3^{(1)}(\mathbf{q},\mathbf{q}_2,\mathbf{q}_1) = \\ &= S_3^{(1)}(\mathbf{q},\mathbf{q}_1,-\mathbf{q}-\mathbf{q}_1-\mathbf{q}_2) \\ &= S_3^{(1)}(\mathbf{q},-\mathbf{q}-\mathbf{q}_1-\mathbf{q}_2,\mathbf{q}_2), \end{aligned} \tag{37}$$

and the analogous ones can be written for higher $S_j^{(1)}$.

The equations for the functions $S_j^{(1)}$ can be found from Eq. (30), if we substitute all functions in it and collect all terms referred to each collection in (19) and the constant. In view of the functional difference of collections in (19), we equate these groups of terms to zero. This yields the chain of equations for E_0 and the functions $S_j^{(1)}$:

$$E_0 = \tilde{E}_0^b + A_1, \tag{38}$$

$$\begin{aligned} A_1 \ &= \frac{\hbar^2}{2m}\Bigg[k_1^2 - \frac{1}{N}\sum_{\mathbf{q}\neq 0}^{(2\pi)} q^2 S_1^{(1)}(\mathbf{q})S_1^{(1)}(-\mathbf{q}) - \\ &- \ \frac{i}{\sqrt{N}}\sum_{q_x\neq 0}^{(2\pi)} 2k_{1x}q_x C_{1x}(q_x)S_1^{(1)}(-\mathbf{q}_x) + (x \to y, z)\Bigg], \end{aligned} \tag{39}$$

$$
\begin{aligned}
&S_1^{(1)}(\mathbf{q})\epsilon_0(\mathbf{q}) = \\
= \;& -i\sqrt{N}2k_{1x}q_x a_2(-\mathbf{q}_x)C_{1x}(q_x)\delta_{\mathbf{q},\mathbf{q}_x} + (x \to y,z) + \\
+ \;& \frac{2}{N}\sum_{\mathbf{q}_1\neq 0}^{(\pi)} \Big\{(q_1^2+\mathbf{q}_1\mathbf{q})S_2^{(1)}(\mathbf{q},\mathbf{q}_1) + \\
+ \;& 3q_1^2 S_3^{(1)}(\mathbf{q},\mathbf{q}_1,-\mathbf{q}_1) + \\
+ \;& \sqrt{N}2q_1^2 S_1^{(1)}(\mathbf{q}_1)S_2^{(1)}(\mathbf{q}-\mathbf{q}_1,\mathbf{q}_1) + \\
+ \;& 0.5\sqrt{N}(q_1^2-\mathbf{q}_1\mathbf{q})S_1^{(1)}(\mathbf{q}_1)S_1^{(1)}(\mathbf{q}-\mathbf{q}_1)\Big\} + \\
+ \;& \left\{\sum_{p_x\neq 0}^{(2\pi)} 2ik_{1x}C_{1x}(p_x)\Big[(q_x-p_x)S_1^{(1)}(\mathbf{q}-\mathbf{p}_x) - \right. \\
- \;& \left. 2p_x S_2^{(1)}(\mathbf{q}-\mathbf{p}_x,-\mathbf{q})\Big] + (x\to y,z)\right\}, \qquad (40)
\end{aligned}
$$

$$
\begin{aligned}
&S_2^{(1)}(\mathbf{q},\mathbf{q}_1)\left[\epsilon_0(\mathbf{q}_1)+\epsilon_0(\mathbf{q}+\mathbf{q}_1)\right] + \\
+ \;& 2S_1^{(1)}(\mathbf{q})a_2(-\mathbf{q}_1)\mathbf{q}\mathbf{q}_1 - q^2 S_1^{(1)}(\mathbf{q})a_3(\mathbf{q},\mathbf{q}_1) = \\
= \;& \delta_{\mathbf{q},\mathbf{q}_x}\sqrt{N}ik_{1x}C_{1x}(q_x)\left\{2q_{1x}a_2(\mathbf{q}_1) - q_x a_3(\mathbf{q}_1,\mathbf{q}_x)\right\} + \\
+ \;& \sum_{p_x\neq 0}^{(2\pi)} ik_{1x}C_{1x}(p_x)\Big[4(q_x+q_{1x}-p_x)S_2^{(1)}(\mathbf{q}-\mathbf{p}_x,\mathbf{q}_1) - \\
- \;& 6p_x S_3^{(1)}(\mathbf{q}-\mathbf{p}_x,\mathbf{q}_1,-\mathbf{q}-\mathbf{q}_1)\Big] + (x\to y,z) + \\
+ \;& \frac{1}{\sqrt{N}}\sum_{\mathbf{q}_2\neq 0}^{(\pi)}\Big\{\frac{2}{\sqrt{N}}(q_2^2-\mathbf{q}_1\mathbf{q}_2)S_3^{(1)}(\mathbf{q},\mathbf{q}_1-\mathbf{q}_2,\mathbf{q}_2) + \\
+ \;& \frac{4}{\sqrt{N}}\mathbf{q}_2(\mathbf{q}_1+\mathbf{q}_2+\mathbf{q})S_3^{(1)}(\mathbf{q},\mathbf{q}_1,\mathbf{q}_2) - \\
- \;& 4\mathbf{q}_2(\mathbf{q}_1-\mathbf{q}_2+\mathbf{q})S_1^{(1)}(\mathbf{q}_2)S_2^{(1)}(\mathbf{q}-\mathbf{q}_2,\mathbf{q}_1) + \\
+ \;& 6q_2^2 S_1^{(1)}(\mathbf{q}_2)S_3^{(1)}(\mathbf{q}-\mathbf{q}_2,\mathbf{q}_1,-\mathbf{q}-\mathbf{q}_1) + \\
+ \;& 4(\mathbf{q}_1+\mathbf{q}_2)^2 S_2^{(1)}(\mathbf{q}_2,\mathbf{q}_1)S_2^{(1)}(\mathbf{q}-\mathbf{q}_2,\mathbf{q}_1+\mathbf{q}_2)\Big\}. \qquad (41)
\end{aligned}
$$

For brevity, we do not give the equations for the higher functions $S_{j\geq 3}^{(1)}(\mathbf{q})$. Equation (41) should be symmetrized to satisfy (36). For the validity of relation (30), it is sufficient to consider only $S_j^{(1)}$ with the first even argument. We have the rule: the first argument of the functions $S_j^{(1)}$ is always quantized by the even law (2). Otherwise, we set $S_j^{(1)} = 0$.

Analogously, (29) yields the chain of equations for $\tilde{E}_0^b$ and the functions a_j:

$$
\tilde{E}_0^b = \frac{N-1}{2^{d+1}N}n\nu_3(0) - \sum_{\mathbf{k}\neq 0}^{(\pi)}\frac{n\nu_3(k)}{2^{d+1}N} - \sum_{\mathbf{k}\neq 0}^{(\pi)}\frac{\hbar^2 k^2}{2mN}a_2(\mathbf{k}), \qquad (42)
$$

$$
\begin{aligned}
&\frac{n\nu_3(k)m}{2^d\hbar^2} + a_2(\mathbf{k})k^2 - a_2^2(\mathbf{k})k^2 = A_2(\mathbf{k}) + \qquad (43) \\
&+ \sum_{\mathbf{q}\neq 0}^{(\pi)}\frac{a_3(\mathbf{k},\mathbf{q})}{N}(q^2+\mathbf{k}\mathbf{q}) + \sum_{\mathbf{q}\neq 0}^{(\pi)} q^2\frac{a_4(\mathbf{q},-\mathbf{q},\mathbf{k})}{2N},
\end{aligned}
$$

$$
a_3(\mathbf{k},\mathbf{q}) \approx A_3(\mathbf{k},\mathbf{q}) - \frac{2R(\mathbf{k},\mathbf{q})}{\epsilon_0(k)+\epsilon_0(q)+\epsilon_0(\mathbf{k}+\mathbf{q})}. \qquad (44)
$$

These equations can be found from (6)–(9) by the changes $\nu_3(k) \to \nu_3(k)/2^d$, $\sum\limits^{(2\pi)} \to \sum\limits^{(\pi)}$ and the addition of “surface” corrections A_j. The latter are related to the appearance of the terms

$$
-\frac{\hbar^2}{2m}\sum_{\mathbf{k}\neq 0}^{(\pi)} A_2(\mathbf{k})\rho_{\mathbf{k}}\rho_{-\mathbf{k}},
$$

$$
-\frac{\hbar^2}{2m}\sum_{\mathbf{k}_1,\mathbf{k}_2\neq 0}^{(\pi)\mathbf{k}_1+\mathbf{k}_2\neq 0} A_3(\mathbf{k}_1,\mathbf{k}_2)\rho_{\mathbf{k}_1}\rho_{\mathbf{k}_2}\rho_{-\mathbf{k}_1-\mathbf{k}_2}, \ldots
$$

in Eq. (30), which have the structure of terms of Eq. (29). In particular,

$$
\begin{aligned}
A_2(\mathbf{k}) \;=\;& \sum_{q_x\neq 0}^{(2\pi)}\frac{ik_{1x}C_{1x}(-q_x)}{\sqrt{N}}\Big[4(q_x+k_x)S_2^{(1)}(\mathbf{q}_x,\mathbf{k}) + \\
+ \;& 6q_x S_3^{(1)}(\mathbf{q}_x,-\mathbf{q}_x,\mathbf{k})\Big] + (x\to y,z) + \\
+ \;& \frac{1}{N}\sum_{\mathbf{q}\neq 0}^{(2\pi)}\Big[4(q^2+\mathbf{q}\mathbf{k})S_1^{(1)}(-\mathbf{q})S_2^{(1)}(\mathbf{q},\mathbf{k}) + \\
+ \;& 4(\mathbf{k}+\mathbf{q})^2 S_2^{(1)}(\mathbf{q},\mathbf{k})S_2^{(1)}(-\mathbf{q},-\mathbf{k}) + \\
+ \;& 6q^2 S_1^{(1)}(-\mathbf{q})S_3^{(1)}(\mathbf{q},\mathbf{k},-\mathbf{k})\Big]. \qquad (45)
\end{aligned}
$$

Though the majority of terms in (29) and (30) has the structure of the collections $\rho_{\mathbf{k}}$, which is inherent only in the given equation ((29) or (30)), a small part of terms in (30) has the structure of terms in (29). The criterion for the construction of the chains of equations for $S_j^{(1)}$ and a_j is the functional difference of the collections $\rho_{\mathbf{k}}$ (19). Basically, the structures of the terms in (29) and (30) are different. Therefore, the partition of Eq. (18) into (29) and (30) is justified and simplifies the determination of $S_w^{(1)}$. The collections $\rho_{\mathbf{k}}$, which are present in both equations, should be analyzed with the use of the initial equation (18). We can verify that Ψ_0 (26) satisfies Eq. (18), if a_j and A_j satisfy Eqs. (42)–(45).

As $U_s \to \infty$ and $\delta \to 0$, Eqs. (26), (35), (38)–(45) are the *required solution* for the WF of the ground state of N interacting Bose particles filling a rectangular vessel.

The analysis of Eqs. (40) and (41) indicates that the functions $S_1^{(1)}(\mathbf{q})$ and $S_2^{(1)}(\mathbf{q},\mathbf{q}_1)$ can be considered “one-dimensional”:

$$
S_1^{(1)}(\mathbf{q}) \approx S_1^{(1)}(\mathbf{q})(\delta_{\mathbf{q},\mathbf{q}_x}+\delta_{\mathbf{q},\mathbf{q}_y}+\delta_{\mathbf{q},\mathbf{q}_z}), \qquad (46)
$$

$$
\begin{aligned}
S_2^{(1)}(\mathbf{q},\mathbf{q}_1) \;&\approx \delta_{\mathbf{q},\mathbf{q}_x}S_2^{(1)}(\mathbf{q}_x,\mathbf{q}_1) + \\
+ \;&\delta_{\mathbf{q},\mathbf{q}_y}S_2^{(1)}(\mathbf{q}_y,\mathbf{q}_1) + \delta_{\mathbf{q},\mathbf{q}_z}S_2^{(1)}(\mathbf{q}_z,\mathbf{q}_1) \qquad (47)
\end{aligned}
$$

($\mathbf{q} \equiv (q_x, q_y, q_z) = \mathbf{q}_x + \mathbf{q}_y + \mathbf{q}_z$), since the nonone-dimensional $S_j^{(1)}$ with $\mathbf{q} = (q_x, q_y, 0)$ (with possible permutations of the x-, y-, and z-components) and $\mathbf{q} = (q_x, q_y, q_z)$ are less than one-dimensional ones by, respectively, $N^{1/3}$ and $N^{2/3}$ times. Due to the smallness of nonone-dimensional terms, their inclusion in the equations does not influence the values of one-dimensional ones and the value of E_0. For the one-dimensional parts of the functions $S_j^{(1)}$, all terms in Eqs. (40) and (41) are of the same order, as it is easy to verify. Therefore, all they should be taken into account. But, in this case, (38)–(41) is a complicated system of nonlinear integral equations difficult to be solved. Below, we will use the simplest zero approximations for $S_1^{(1)}$ and $S_2^{(1)}$:

$$S_1^{(1)}(\mathbf{q}_x) \approx -\frac{i\sqrt{N} k_{1x} C_{1x}(q_x) 2a_2(-\mathbf{q}_x)}{q_x - 2q_x a_2(\mathbf{q}_x)}, \tag{48}$$

$$S_2^{(1)}(\mathbf{q}_x, \mathbf{q}_1) \approx \frac{i\sqrt{N} k_{1x} C_{1x}(q_x)}{1 - 2a_2(\mathbf{q}_x)} \frac{2q_{1x} a_2(\mathbf{q}_1) - q_x a_3(\mathbf{q}_1, \mathbf{q}_x)}{\epsilon_0(\mathbf{q}_1) + \epsilon_0(\mathbf{q}_x + \mathbf{q}_1)}. \tag{49}$$

Relations (34), (48), and (49) yield $S_1^{(1)}(\mathbf{q}_x) \sim (k_{rot}/q_x) N^{1/6}$ and $S_2^{(1)}(\mathbf{q}_x, \mathbf{q}_1) \sim N^{1/6} \frac{q_x}{q_1 + q_x} (\frac{q_{1x}}{q_1} + \frac{q_x}{q_1 + q_x})$ at $q_x \lesssim k_{rot}$ (k_{rot} is the roton momentum). So, $S_2^{(1)}$ takes values from $\sim N^{-1/2}$ to $\sim N^{1/6}$.

It is easy to verify that the surface corrections A_1 and A_2 have no effect on the solutions for E_0 and $a_2(k)$: the values of A_1 and A_2 by $N^{1/3}$ times less than those, which would render some effect. We expect that $A_{j \geq 3}$ do not affect the solutions for $a_{j \geq 3}$. However, the boundaries change the "bulk part" of the equations for E_0 and a_j: the network of vectors $\mathbf{k}$ becomes by a factor of 2^d denser, and the Fourier-transform of the potential is multiplied by $1/2^d$. For this reason, the boundaries strongly influence the values of E_0 and a_j.

Besides WF (26), WFs of the form (26) with $\Psi_{\mathbf{0}}^f \to \Psi_{\mathbf{l}}^f$ (and $\mathbf{k}_1 \to \mathbf{k}_{\mathbf{l}}$) are also the solutions. But such WFs are equal to zero inside the vessel, whereas Ψ_0 (26) is nonzero. The theorem on nodes implies that the state with a WF without nodes corresponds to the minimal energy. Therefore, the quantity Ψ_0 is the exact and unique WF for the ground state of a Bose liquid and Bose gas in a box.

III. GROUND STATE OF HE II: SOLUTIONS

To verify the solution (26), we will find the behavior of Ψ_0 in the case where the coordinate x of one of the atoms, for example the first one, varies from 0 to L_x. The sines standing before the exponential function in (26) set an inhomogeneous arc-like distribution of atoms in the vessel. But the density of He II is constant in the whole volume of the vessel except for a narrow strip ($\sim$ Å) near the wall. The behavior of Ψ_0 should be analogous. Since the function $e^{\tilde{S}_b}$ sets a homogeneous distribution of atoms, we need to consider the behavior of the ansatz $\Psi_{\mathbf{0}}^f e^{S_w^{(1)}}$ in (26). The function $S_w^{(1)}(\mathbf{r}_1, \ldots, \mathbf{r}_N)$ (35) contains the one-particle dependence in each sum, and the values are of the same order. But taking account of the sums with $S_{j \geq 2}^{(1)}$ is a complicated task, and we neglect it. Then

$$\Psi_0(x_1) \sim f(\mathbf{r}_1) = \sin(k_{1x} x_1 + \delta_{1x}) e^{s_1(\mathbf{r}_1)}, \tag{50}$$

$$\begin{aligned} s_1(\mathbf{r}_1) = & \sum_{q_x \neq 0}^{(2\pi)} \frac{S_1^{(1)}(\mathbf{q}_x)}{\sqrt{N}} e^{iq_x x_1} + \\ + & \sum_{q_x, q_y \neq 0}^{(2\pi)} \frac{S_1^{(1)}(\mathbf{q}_x + \mathbf{q}_y)}{\sqrt{N}} e^{iq_x x_1 + iq_y y_1} + \\ + & \sum_{q_x, q_z \neq 0}^{(2\pi)} \frac{S_1^{(1)}(\mathbf{q}_x + \mathbf{q}_z)}{\sqrt{N}} e^{iq_x x_1 + iq_z z_1} + \\ + & \sum_{q_x, q_y, q_z \neq 0}^{(2\pi)} \frac{S_1^{(1)}(\mathbf{q}_x + \mathbf{q}_y + \mathbf{q}_z)}{\sqrt{N}} e^{iq_x x_1 + iq_y y_1 + iq_z z_1}. \end{aligned} \tag{51}$$

As was noted above, we have $S_1^{(1)}(\mathbf{q}_x + \mathbf{q}_y + \mathbf{q}_z) \sim S_1^{(1)}(\mathbf{q}_x + \mathbf{q}_y)/N^{1/3} \sim S_1^{(1)}(\mathbf{q}_x)/N^{2/3}$. Therefore, three last sums in (51) are of the order of the first one, but they depend additionally on y_1 and z_1. Let us consider only the first sum:

$$s_1(\mathbf{r}_1) = s_1(x_1) = \sum_{q_x = 2\pi j_x/L_x}^{(2\pi)} \frac{S_1^{(1)}(\mathbf{q}_x)}{\sqrt{N}} e^{iq_x x_1}, \tag{52}$$

where $j_x = \pm 1, \pm 2, \pm 3, \ldots$. For $S_1^{(1)}(\mathbf{q}_x)$, we use the zero approximation (48). In view of the property $S_1^{(1)}(-\mathbf{q}_x) = S_1^{(1)}(\mathbf{q}_x)$ and since $S_1^{(1)}(\mathbf{q}_x)$ is real, the imaginary part of (52) is cancelled, and we can change $e^{iq_x x_1} \to \cos(q_x x_1)$. For investigating the qualitative behavior of the solutions, we choose the interatomic potential of two He^4 atoms in the simple form

$$U_3(\mathbf{r}) \approx \left[\begin{array}{cc} U_b > 0 & r \leq a \\ U_{bd} < 0 & a \leq r \leq b \\ 0 & r > b. \end{array} \right. \tag{53}$$

For the modern potentials [33, 34], $a \approx 2.6$ Å, $U_{bd} = -11$ K, but the barrier and the well are more flat than the steps (53). Therefore, we will use in (53) the lower values of a and $|U_{bd}|$: $a = 2$ Å, $U_{bd} = -9$ K, and $b = 4$ Å.

The Fourier-transform of potential (53) is

$$\nu_3(k) = \frac{4\pi}{k^3} \tilde{\nu}_3(k), \tag{54}$$

$$\begin{aligned} \tilde{\nu}_3(k) & = [U_b - U_{bd}][\sin(ak) - ak\cos(ak)] + \\ & + U_{bd}[\sin(bk) - bk\cos(bk)]. \end{aligned} \tag{55}$$

In the two-particle approximation, we have from (43):

$$\begin{aligned} a_2(k) &= \frac{1}{2} - \sqrt{\frac{1}{4} + \frac{n\nu_3(k)m}{2^d\hbar^2k^2}} = \\ &= \frac{1}{2} - \sqrt{\frac{1}{4} + \frac{\alpha_1\tilde{\nu}_3(k)}{(\bar{R}k)^5}}, \qquad (56) \end{aligned}$$

where $\alpha_1 = 4\pi m\bar{R}^2/2^d\hbar^2 \approx 1.66/Kk_B$, $d = 3$, and $\bar{R} \approx 3.58\,\text{Å}$ is the mean interatomic distance for He II. For a_2, we chose the sign "minus" before the root. At the sign "plus," we have nonphysical $E_0 = -\infty$.



Fig.1. Values of the functions $F = f(x)$ (○) and $F = s_1(x)$ (⋆) by formulas (50), (52) for potentials (14) and (53) at $a = 2\,\text{Å}$, $U_b = 1600\,\text{K}$, $U_{bd} = 0\,\text{K}$, $U_s \to \infty$ and the vessel size $L_x = 10^4\bar{R}$. Values of $f(x)$ and $s_1(x)$ at $x/L_x \in [0.5, 1]$ are obtained by the reflection of the curves in an upright mirror, positioned at $x = 0.5L_x$.

The results for the functions $f(x_1)$ and $s_1(x_1)$ (50), (52) at $L_x = 10^4\bar{R}$ and several potentials are given in Figs. 1 and 2. The values of the functions at $L_x = 10^5\bar{R}$ and $L_x = 10^3\bar{R}$ differ from the drawn ones very slightly (by $\lesssim 0.1\%$). So, the curves in the figures are true for all $L_x \gtrsim 10^3\bar{R}$. As seen from Fig. 1, $f(x_1)$ is equal to 0.5 everywhere, except for a narrow region near the wall. While approaching the wall, $f(x_1)$ oscillates with constant period and increasing amplitude and becomes zero on the wall. The dependence on the potential is as follows. An increase of the interatomic barrier U_b causes the growth of the amplitude of oscillations $f(x_1)$. The well depth U_{bd} does not affect the results, if $|U_{bd}| \ll U_b$. As the distance to the wall increases, $f(x_1)$ approaches the constant: $f(x_1) = 0.5$. This holds with a high accuracy increasing with L_x. For example, at $L_x = 10^4\bar{R}$, the deviation from 0.5 occurs only in the sixth decimal point. As $x_1 \to L_x$, the function $f(x_1)$ behaves itself as at $x_1 \to 0$.



Fig. 2. Values of the function $f(x)$ (50), (52) near the wall ($x = 0$) at $L_x = 10^4\bar{R}$, $U_s \to \infty$ and potential (53) with $a = 2\,\text{Å}$, $b = 4\,\text{Å}$, $U_{bd} = -9\,\text{K}$, and three barriers: $U_b = 800\,\text{K}$ (○), $U_b = 2000\,\text{K}$ (⋆), and $U_b = 2900\,\text{K}$ (◇). At $x \gtrsim 6\,\text{Å}$, curves ○ ○ ○ and ⋆ ⋆ ⋆ almost merge. Values of x are given in Å.

It is of interest that $f(x_1)$ oscillates near the wall. This is because the short-range order, and the period is proportional to $a/\bar{R}$. Large oscillations at $U_b = 2900\,\text{K}$ are, most likely, nonphysical and are related to the neglect of corrections, which are large at such a potential. This reflects the behavior of the density ρ of helium, since $\rho(x_1) \sim \int dy_1dz_1d\mathbf{r}_2\ldots d\mathbf{r}_N|\Psi_0|^2 = const \times f^2(x_1)$ (the behavior of $\rho(x_1)$ near the wall is also affected by the integral of $e^{\tilde{S}_b}$, but we did not study it). Similar oscillations were obtained in [21], while modeling the properties of a film of helium adsorbed on the substrate.

We obtained the solutions also at finite U_s. At $U_s \gtrsim 1000\,\text{K}$, the solutions are practically invariable with increase in U_s, except for the values on a wall, where $f \neq 0$. As U_s increases, the function $s_1(x)$ approaches a constant ($\simeq 10$) at the point $x = 0$, and $\sin(k_{1x}x_1 + \delta_{1x}) \to 0$. As a result, we have $f(x \to 0)|_{U_s\to\infty} \to 0$. With regard for the higher corrections, such a behavior will be conserved: as $U_s \to \infty$, we have $\delta \to 0$, such δ affect slightly $C_1(q)$ and all results, because δ enter the equations through the function $C_1(q)$. In addition, the terms with large q are cut in sums by a_j. Due to such properties, the transition to opaque walls ($U_s \to \infty$, $\delta \to 0$) is correct and makes Ψ_0 to be arbitrarily close to zero on the walls.

Thus, though the bare function $\sin(k_{1x}x_1 + \delta_{1x})$ is far from to be a constant, the account for $S_1^{(1)}(q_x)$ makes the theoretical density of atoms to be constant everywhere, except for a narrow ($\sim$ Å) region near walls. Therefore, $\sin(k_{1x}x_1 + \delta_{1x})$ is a "good" bare function, despite the anxiety [9–11]. It satisfies the equation for free particles. If the interatomic interaction is "switched-off," Ψ_0 is reduced to product (20) of such functions, as it should be. Apparently, this is the best bare function: at the other bare functions, we obtain the same solution for Ψ_0, but in other form. However, the properties of the solution are not improved in this case, whereas the equations for $S_j^{(1)}$ become too complicated. The circumstance that the correction $S_1^{(1)}(q_x)$ "improve" the behavior of $f(x_1)$ seems natural, because the Schrödinger equation (18) involves the interaction of atoms and must make their distribution uniform, but $S_1^{(1)}(q_x)$ is generated just by the Schrödinger equation.

We can find E_0 (38), (42) with regard for $A_1 \approx 0$ and a_2 (56). However, the potential of a semitransparent ball (53) is "bad" for this purpose in view of the asymptotics $\nu_3(k \to \infty) \sim \cos{(ak)}/k^2$, at which the value of E_0 (42), (56) is uncertain. For 3D-systems, the suitable asimptotics is $\nu_3(k \to \infty) \sim 1/k^p, p > 3$. Therefore, we will not estimate E_0 here.

IV. EXCITED STATE OF HE II

We now find the WF of helium II in a vessel for the state with one phonon. Under periodic BCs, the exact solution for the WF takes the form [27]

$$\Psi_{\mathbf{k}}(\mathbf{r}_1, \ldots, \mathbf{r}_N) = \psi_{\mathbf{k}}\Psi_0, \tag{57}$$

$$\psi_{\mathbf{k}} \equiv \psi^b_{\mathbf{k}} = \rho_{-\mathbf{k}} + \sum^{(2\pi)}_{\mathbf{k}_2 \neq 0, -\mathbf{k}} \frac{b_2(\mathbf{k}_2, \mathbf{k})}{\sqrt{N}} \rho_{\mathbf{k}_2}\rho_{-\mathbf{k}_2-\mathbf{k}} + \tag{58}$$
$$+ \sum^{(2\pi)\, \mathbf{k}_2+\mathbf{k}_3+\mathbf{k} \neq 0}_{\mathbf{k}_2, \mathbf{k}_3 \neq 0} \frac{b_3(\mathbf{k}_2, \mathbf{k}_3, \mathbf{k})}{N} \rho_{\mathbf{k}_2}\rho_{\mathbf{k}_3}\rho_{-\mathbf{k}_2-\mathbf{k}_3-\mathbf{k}} + \ldots$$

The approximate $\psi_{\mathbf{k}}$ of such a type, which consists of the zero approximation

$$\psi^b_{\mathbf{k}} = \rho_{-\mathbf{k}} \tag{59}$$

and a simplified correction with b_2, was first proposed by R. Feynman [4, 5]. The arguments to favor of solution (57), (58) are as follows: 1) it satisfies the Schrödinger equation for N interacting Bose particles and 2) it is an eigenfunction of the momentum operator $\hat{\mathbf{P}} = -i\hbar \sum\limits_j \nabla_{\mathbf{r}_j}$ with the eigenvalue $\hbar\mathbf{k}$; 3) at the "switching-off" of the interaction, it is reduced to WF (59) describing N free Bose particles, from which $N-1$ particles are immovable, and a single one moves with the momentum $\mathbf{k}$; 4) an analogous solution is obtained in the operator approach [3]; 5) the completeness [17] of collections (19) implies that the solution for $\psi_{\mathbf{k}}$ should have only such a form. However, the structure of (57) is only the assumption, though no solution with another structure was found. The solution for $\psi_{\mathbf{k}}$ with "shadow" variables [22, 23] is equivalent to (57), (58).

For the fluid in a vessel, the WF $\Psi_{\mathbf{k}}$ should be sought anew, since $\Psi_{\mathbf{k}}$ (57), (58) is not the solution of the Schrödinger equation.

The simplest way to find solution (57), (58) is to assume the structure of (57) and to substitute the WF of free particles (59), as the bare $\psi_{\mathbf{k}}$, in the Schrödinger equation for interacting particles. The equation prompts the form of corrections to $\psi_{\mathbf{k}}$.

The reasoning for particles in a vessel is analogous. We start from (57) with Ψ_0 (26). The study of different possibilities has shown that $\psi_{\mathbf{k}}$ must be sought in the form

$$\psi_{\mathbf{k}} = \frac{8}{\sqrt{N}} \sum_{j=1}^{N} \cos{(k_{r_x} x_j)} \cos{(k_{r_y} y_j)} \cos{(k_{r_z} z_j)} + \delta\psi_{\mathbf{k}} \tag{60}$$

with the unknown $\delta\psi_{\mathbf{k}}$. In this case, $\mathbf{k_r} = (k_{r_x}, k_{r_y}, k_{r_z})$ takes values from series (3), since we must obtain the solution for free particles in a box with law (3) at the switching-off of the interaction. We note that

$$\frac{8}{\sqrt{N}} \sum_{j=1}^{N} \cos{(k_{r_x} x_j)} \cos{(k_{r_y} y_j)} \cos{(k_{r_z} z_j)} =$$
$$= \rho_{-\mathbf{k_r}} + 7\,\text{permutations}, \tag{61}$$

where the permutation means $\rho_{-\mathbf{k}}$ with the different sign of one or several components of the vector $\mathbf{k_r}$. We now represent $\psi_{\mathbf{k}}$ (60) in the form

$$\psi_{\mathbf{k}} = \psi^0_{\mathbf{k}} + 7\,\text{permutations}, \tag{62}$$

$$\psi^0_{\mathbf{k}} = \rho_{-\mathbf{k}} + \delta\psi^0_{\mathbf{k}}. \tag{63}$$

It is easy to see that solution (62), (63) describes a three-dimensional standing wave decaying into eight counter running waves.

Substituting WF (57) in the Schrödinger equation with Hamiltonian (12), we obtain the equation for $\psi_{\mathbf{k}}$:

$$-\frac{\hbar^2}{2m} \sum_j \{\triangle_j \psi_{\mathbf{k}} + 2(\nabla_j \psi_{\mathbf{k}}) \times \nabla_j S\} = E(\mathbf{k})\psi_{\mathbf{k}}, \tag{64}$$

$$S = S^{(1)}_w + \tilde{S}_b + S_0, \tag{65}$$

$$S_0 = \ln \Psi^f_{\mathbf{0}}, \tag{66}$$

where $E(\mathbf{k}) = E - NE_0$ is the energy of a quasiparticle. $E(\mathbf{k})$ is obviously independent of the signs of components of $\mathbf{k}$. Therefore, Eq. (64) is separated into 8 equations: for $\psi^0_{\mathbf{k}}$ and 7 permutations with the same energy $E(k)$.

The analysis indicates that the solution for $\psi^0_{\mathbf{k}}$ has the following form:

$$\psi^0_{\mathbf{k}} = \psi^b_{\mathbf{k}} + b_0(\mathbf{k}) + \sum^{(2\pi)}_{\mathbf{q} \neq 0, -\mathbf{k}} Q_1(\mathbf{q}, \mathbf{k})\rho_{-\mathbf{q}-\mathbf{k}} +$$
$$+ \sum^{\mathbf{q}+\mathbf{q}_1+\mathbf{k} \neq 0}_{\mathbf{q}, \mathbf{q}_1 \neq 0} \frac{Q_2(\mathbf{q}, \mathbf{q}_1, \mathbf{k})}{\sqrt{N}} \rho_{\mathbf{q}_1}\rho_{-\mathbf{q}-\mathbf{q}_1-\mathbf{k}} + \tag{67}$$
$$+ \sum^{\mathbf{q}+\mathbf{q}_1+\mathbf{q}_2+\mathbf{k} \neq 0}_{\mathbf{q}, \mathbf{q}_1, \mathbf{q}_2 \neq 0} \frac{Q_3(\mathbf{q}, \mathbf{q}_1, \mathbf{q}_2, \mathbf{k})}{N} \rho_{\mathbf{q}_1}\rho_{\mathbf{q}_2}\rho_{-\mathbf{q}-\mathbf{q}_1-\mathbf{q}_2-\mathbf{k}} +$$
$$+ \ldots,$$

where $\psi^{b}_{\mathbf{k}}$ is the solution (58) for periodic BCs (but with $\mathbf{k}_j$ running series (3)), b_0 and Q_l are the corrections from boundaries. Here, $\mathbf{q}$ is quantized according to (2) (like $2\pi j/L$), and $\mathbf{q}_j$ and $\mathbf{k}$ are quantized by (3) (like $\pi j/L$).

With regard for the properties of $\rho_{\mathbf{k}}$, we can directly verify that $\Psi_{\mathbf{k}}$ (57), (62), (67) is the exact solution of the Schrödinger equation, if the functions $E(k)$, b_j, and Q_l satisfy the equations

$$\begin{aligned}
&b_0(\mathbf{k})\epsilon(k) = -2k^2 S^{(1)}_1(-\mathbf{k})\delta_{\mathbf{k},\mathbf{k}^e} - \\
-\ &2k_{1x}k_x iC_{1x}(-k_x)\sqrt{N}\delta_{\mathbf{k},\mathbf{k}^e_x} - \\
-\ &\frac{2}{\sqrt{N}}\sum^{(\pi)}_{\mathbf{q}\neq 0} q^2 Q_2(-\mathbf{k},\mathbf{q},\mathbf{k})\delta_{\mathbf{k},\mathbf{k}^e} - \\
-\ &\sum^{(2\pi)}_{\mathbf{q}\neq 0,-\mathbf{k}} 2(\mathbf{k}+\mathbf{q})^2 Q_1(\mathbf{q},\mathbf{k})S^{(1)}_1(-\mathbf{k}-\mathbf{q})\delta_{\mathbf{k},\mathbf{k}^e} + \\
+\ &\sum^{(2\pi)}_{q_x\neq 0} 2\sqrt{N}k_{1x}q_x iC_{1x}(q_x)Q_1(-\mathbf{k}-\mathbf{q}_x,\mathbf{k})\delta_{\mathbf{k},\mathbf{k}^e} + \\
+\ &(x\rightarrow y,z), \qquad (68)
\end{aligned}$$

$$\begin{aligned}
\epsilon(k)\ &= \epsilon_0(k) - \frac{1}{N}\sum^{(\pi)}_{\mathbf{k}_2\neq 0} b_2(\mathbf{k}_2,\mathbf{k})2\mathbf{k}_2(\mathbf{k}_2+\mathbf{k}) - \\
-\ &\frac{1}{N}\sum^{(\pi)}_{\mathbf{k}_2\neq 0} 6k_2^2 b_3(\mathbf{k}_2,-\mathbf{k}_2,\mathbf{k}) - \\
-\ &\sum^{(2\pi)}_{q_x\neq 0} 2k_{1x}(k_x+q_x)iC_{1x}(-q_x)Q_1(\mathbf{q}_x,\mathbf{k}) - \\
-\ &\frac{2}{\sqrt{N}}\sum^{(2\pi)}_{\mathbf{q}\neq 0} Q_1(\mathbf{q},\mathbf{k})(\mathbf{k}+\mathbf{q}) \times \\
\times\ &\Big[\mathbf{q}S^{(1)}_1(-\mathbf{q}) + 2(\mathbf{k}+\mathbf{q})S^{(1)}_2(-\mathbf{q},-\mathbf{k})\Big] + \\
+\ &\sum^{(2\pi)}_{q_x\neq 0} 4k_{1x}q_x iC_{1x}(q_x)Q_2(-\mathbf{q}_x,\mathbf{q}_x,\mathbf{k}) - \qquad (69) \\
-\ &\frac{4}{\sqrt{N}}\sum^{(2\pi)}_{\mathbf{q}\neq 0} q^2 Q_2(-\mathbf{q},\mathbf{q},\mathbf{k})S^{(1)}_1(\mathbf{q}) + (x\rightarrow y,z),
\end{aligned}$$

$$\begin{aligned}
&Q_1(\mathbf{q},\mathbf{k})\left[\epsilon(k)-\epsilon_0(\mathbf{k}+\mathbf{q})\right] = \\
=\ &2ik_{1x}C_{1x}(q_x)\delta_{\mathbf{q},\mathbf{q}_x}\left[-k_x + 2q_x b_2(\mathbf{q},\mathbf{k})\right] + \\
+\ &\frac{2S^{(1)}_1(\mathbf{q})}{\sqrt{N}}[\mathbf{q}\mathbf{k} - 2q^2 b_2(\mathbf{q},\mathbf{k})] - \frac{4k^2}{\sqrt{N}}S^{(\mathbf{1})}_2(\mathbf{q},-\mathbf{q}-\mathbf{k}) - \\
-\ &\frac{1}{N}\sum^{(\pi)}_{\mathbf{q}_1\neq 0}\{2\mathbf{q}_1(\mathbf{q}_1+\mathbf{q}+\mathbf{k})\ \times \\
\times\ &[Q_2(\mathbf{q},\mathbf{q}_1,\mathbf{k}) + Q_1(\mathbf{q}+\mathbf{q}_1,\mathbf{k})\sqrt{N}S^{(1)}_1(-\mathbf{q}_1)] + \\
+\ &6q_1^2 Q_3(\mathbf{q},\mathbf{q}_1,-\mathbf{q}_1,\mathbf{k}) + \\
+\ &4q_1^2 Q_2(\mathbf{q}-\mathbf{q}_1,\mathbf{q}_1,\mathbf{k})\sqrt{N}S^{(1)}_1(\mathbf{q}_1) + Q_1(\mathbf{q}+\mathbf{q}_1,\mathbf{k}) \times \\
\times\ &4(\mathbf{q}_1+\mathbf{q}+\mathbf{k})^2\sqrt{N}S^{(1)}_2(-\mathbf{q}_1,-\mathbf{q}-\mathbf{k})\Big\} \\
+\ &\sum^{(2\pi)}_{p_x\neq 0} 2k_{1x}iC_{1x}(p_x)\left[(-k_x-q_x+p_x)Q_1(\mathbf{q}-\mathbf{p}_x,\mathbf{k})\ + \right. \\
+\ &\left. 2p_x Q_2(\mathbf{q}-\mathbf{p}_x,\mathbf{p}_x,\mathbf{k})\right] + (x\rightarrow y,z), \qquad (70)
\end{aligned}$$

$$\begin{aligned}
&b_2(\mathbf{k}_2,\mathbf{k})\left[\epsilon(k)-\epsilon_0(\mathbf{k}_2)-\epsilon_0(\mathbf{k}+\mathbf{k}_2)\right] = \\
=\ &\mathbf{k}\mathbf{k}_2 a_2(\mathbf{k}_2) - \mathbf{k}(\mathbf{k}+\mathbf{k}_2)a_2(\mathbf{k}+\mathbf{k}_2) - k^2 a_3(\mathbf{k},\mathbf{k}_2) - \\
-\ &\frac{2}{\sqrt{N}}\sum^{(\pi)}_{\mathbf{k}_3\neq 0}\Bigg\{\frac{3\mathbf{k}_3(\mathbf{k}_2+\mathbf{k}_3+\mathbf{k})}{\sqrt{N}}b_3(\mathbf{k}_2,\mathbf{k}_3,\mathbf{k})\ + \\
+\ &Q_1(\mathbf{k}_3,\mathbf{k})(\mathbf{k}_3+\mathbf{k})\ \Big[2(\mathbf{k}_3-\mathbf{k}_2)S^{(1)}_2(-\mathbf{k}_3,\mathbf{k}_2)\ + \\
+\ &3(\mathbf{k}_3+\mathbf{k})S^{(1)}_3(-\mathbf{k}_3,\mathbf{k}_2,-\mathbf{k}_2-\mathbf{k})\Big] + \\
+\ &2Q_2(-\mathbf{k}_3,\mathbf{k}_2+\mathbf{k}_3,\mathbf{k})(\mathbf{k}_2+\mathbf{k}_3)\times \\
\times\ &\Big[\mathbf{k}_3S^{(1)}_1(\mathbf{k}_3) + 2(\mathbf{k}_2+\mathbf{k}_3)S^{(1)}_2(\mathbf{k}_3,\mathbf{k}_2)\Big] + \\
+\ &3k_3^2 S^{(1)}_1(\mathbf{k}_3)Q_3(-\mathbf{k}_3,\mathbf{k}_2,-\mathbf{k}_2-\mathbf{k},\mathbf{k})\Big\} + \\
+\ &\sum^{(2\pi)}_{p_x\neq 0} 2k_{1x}iC_{1x}(p_x)\left[3p_x Q_3(-\mathbf{p}_x,\mathbf{k}_2,-\mathbf{k}_2-\mathbf{k},\mathbf{k})\ + \right. \\
+\ &\left. 2(k_{2x}+p_x)Q_2(-\mathbf{p}_x,\mathbf{k}_2+\mathbf{p}_x,\mathbf{k})\right] + \\
+\ &\text{higher corrections} + (x\rightarrow y,z), \qquad (71)
\end{aligned}$$

where $\epsilon(k) = 2mE(k)/\hbar^2$, $\mathbf{k}^e$ is the wave vector with even components (i.e., they are multiple to $2\pi/L$), $\mathbf{q}_x = q_x\mathbf{i}_x, \mathbf{q}_y = q_y\mathbf{i}_y, \mathbf{q}_z = q_z\mathbf{i}_z$, and analogously for $\mathbf{p}_x$, $\mathbf{p}_y$, and $\mathbf{p}_z$. By $(x\rightarrow y,z)$, we denote the same terms as that with the separated x-component, but with the changes $x\rightarrow y$ and $x\rightarrow z$. All terms of Eqs. (68)–(71), except for the first one, contain a product of two wave vectors (for example, q^2, $k_{1x}k_x$, or $(\mathbf{k}_3+\mathbf{k})(\mathbf{k}_3-\mathbf{k}_2)$). These wave vectors must be nonzero.

Equations (68)–(71) are written in the approximation of “two sums in the wave vector”, at which the series contain the functions a_2, a_3, b_2, b_3, $S^{(1)}_{j\leq 3}$, and $Q_{l\leq 3}$ and do not include the corrections $a_{j\geq 4}$, $b_{j\geq 4}$, $S^{(1)}_{j\geq 4}$, and $Q_{l\geq 4}$. Taking these corrections into account will change the equation for b_2, by adding the correcting sums with b_4, $S^{(1)}_4$, and Q_4.

If the interaction is switched-off, $\Psi_{\mathbf{k}}$ reveals a complicated structure, which should be reduced to the solution for free particles; but we omit this point.

Let us consider the properties of Eqs. (67)–(71). Solution (58) was obtained under periodic BCs and contains only the functions $b_{j\geq 2}$. Solution (67) corresponds to the zero BCs and contains the additional functions b_0 and Q_l. In view of the "one-dimensional" form (46), (47) of the functions $S_j^{(1)}$, relations (68)–(71) yield that the functions Q_l can be also considered "one-dimensional." In particular,

$$Q_1(\mathbf{q},\mathbf{k}) \approx Q_1(\mathbf{q},\mathbf{k})(\delta_{\mathbf{q},\mathbf{q}_x}+\delta_{\mathbf{q},\mathbf{q}_y}+\delta_{\mathbf{q},\mathbf{q}_z}). \quad (72)$$

The "two-dimensional" Q_1 with $\mathbf{q}$ of the form $(q_x, q_y, 0)$ and the "three-dimensional" ones with $\mathbf{q} = (q_x, q_y, q_z)$ are less than the one-dimensional ones by $\sim N^{1/3}$ and $\sim N^{2/3}$ times, respectively. Due to the smallness of nonone-dimensional Q_1, their inclusion in the equations does not affect the values of one-dimensional ones. The estimates indicate that Q_l is less than $k_1 k$ and affects slightly $\epsilon(k)$, since the bulk corrections (the second and third terms on the right-hand side of (69)) are of the order of $\epsilon_0(k) \sim k k_{rot}$, i.e., by $\sim N^{1/3}$ times larger. In Eq. (71), the surface corrections for b_2 are also less than the bulk ones by $\sim N^{1/3}$ times. The smallness of surface corrections A_l and Q_l is obviously understandable and is related to the thinness of a near-surface fluid layer (Fig. 2) as compared with the sizes of a system.

Therefore, we can set $Q_l = 0$ in Eqs. (67)–(71). Then we obtain the well-known Vakarchuk–Yukhnovskii equations [27] for $\epsilon(k)$ and b_j. However, the wave vectors $\mathbf{k}_j$ in these equations should be quantized by the law π/L (3) instead of the usually used $2\pi/L$-law (2), which is valid for the periodic BCs. In the zero approximation for Ψ_0 and $\Psi_{\mathbf{k}}$, we set $a_{j\geq 3} = 0$ and $b_j = 0$ in the equations. Then we obtain the Bogolyubov formula for the energy of quasiparticles,

$$E(k) \approx \sqrt{\left(\frac{\hbar^2 k^2}{2m}\right)^2 + \frac{n\nu_3(k)}{2^d}\frac{\hbar^2 k^2}{m}}, \quad (73)$$

where, however, the potential has the additional factor $1/2^d$ due to effect of the boundaries (in the general case, d is the number of *nonperiodic coordinates*). This formula is true at a weak interaction. Comparing formula (73) at potential (53) with the dispersion curve of He II as $k \to 0$, we obtain $U_b \simeq 283\,\mathrm{K}$, whereas we have $U_b \simeq 35\,\mathrm{K}$ without the factor $1/2^d$. The first estimate is more plausible.

The Bogolyubov-Zubarev field model [3] gives the same results: Since the potential should be expanded in [3] in the Fourier series (15), we replace $\nu_3(k) \to \nu_3(k)/2^d$ and $\overset{(2\pi)}{\sum} \to \overset{(\pi)}{\sum}$ in formulas [3] and obtain the formulas for $E(k)$ (73) and E_0 (42) with $a_2(k)$ (56).

For a Bose gas in a trap, the sound velocity was measured [35]. Its value agrees approximately with the formula $c \approx \sqrt{n\nu_3(0)/2m}$. We cannot compare these results with ours, because our calculations did not involve external fields, whereas the field of a trap plays a significant role in the experiment in [35].

Relations $\partial\langle\hat{H}\rangle/\partial\nu_3(k) = \langle\partial\hat{H}/\partial\nu_3(k)\rangle$, $S(k) = \langle\rho_{\mathbf{k}}\rho_{-\mathbf{k}}\rangle$ and (16) yield [27] $S(k) = 1 + (2^{d+1}/n)(\partial E_0/\partial\nu_3(k))$. Using this result, (38), and (42), we will find the connection between $a_2(k)$ and the structural factor $S(k)$. In the zero approximation, it is the same like that for periodic BCs: $1 - 2a_2(k) \approx 1/S(k)$. With regard for this relation and (56), formula (73) is transformed into the Feynman formula

$$E(k) \approx \hbar^2 k^2/2mS(k), \quad (74)$$

which agrees approximately with the experiment.

If one particle is in the box, k is restricted from above by the value $k^{max} = \sqrt{6mU_s}/\hbar$ (see (25)). For He^4 atoms at $U_s = 25\,K$, we have $k^{max} \approx 3.6\,\mathrm{\AA}^{-1}$. So, the observed break of the dispersion curve of He II at $k \approx 3.6\,\mathrm{\AA}^{-1}$ can be caused, in principle, by that a phonon must have also some k^{max} at a finite barrier of the wall, rather than by the decay mechanism [36].

Additional comments for $\psi_{\mathbf{k}}$ can be found in Appendix A.

V. ROLE OF BOUNDARIES

Let us return to the effect of boundaries, which is delicate. As usual, the works on quantum fluids and crystals do not consider the boundaries and use periodic BCs. The discussion of a role of boundaries can be found only in a few works. In particular, it was noted by J. Ziman [37] that there exists a bulky proof of the correctness of periodic BCs (but it is not presented), and that the identical density of states (for quasiparticles) under real and periodic BCs testifies to the applicability of the latter. K. Huang [11] gave the approximate reasoning in favor of periodic BCs, and expressed the *cautious hope* for that the periodic BCs do not lead to erroneous results.

It was mathematically shown above that this hope is not justified: the boundaries change significantly the solutions. How we can imagine this? According to Sec. 2, BCs reveal themselves at the expansion of the potential $U_3(|\mathbf{r}_1 - \mathbf{r}_2|)$ in a Fourier series: the expansion is realized with wave vectors multiple to $2\pi/L$ under periodic BCs and to π/L under zero BCs. This becomes clear in the following example. Let us consider oscillations of two identical strings with length L: one string is unclosed and has fixed ends, whereas another one is closed. The unclosed string admits the standing waves with the integer number of the waves ($L = j\lambda, j = 1, 2, 3\ldots$) or half-waves ($L = j\lambda/2, j = 1, 3, 5\ldots$) on the string. But, for the closed string, only waves of the first type are realized. Periodic BCs and zero BCs correspond to closed and unclosed strings, respectively. We see that the collections of natural oscillatory harmonics are different in both cases.

The ground state of a Bose fluid can be compared with an oscillating string, since $\Psi_0 \simeq \exp\left(\sum_{\mathbf{k}} a_2(\mathbf{k})\rho_{\mathbf{k}}\rho_{-\mathbf{k}}/2\right)$ has the structure of the WF of the ground state of oscillators with "coordinates" $\rho_{\mathbf{k}}$. The boundaries change the collection of harmonics of the GS. Due to the interaction of harmonics, their intensity is changed as well. Therefore, the solutions for Ψ_0 are different for periodic and zero BCs. Moreover, due to the coupling of the equations for the ground and excited states, the harmonics of the GS affect the excited states. Thus, though the density of excited states is the same for periodic and zero BCs, the sets of $\mathbf{k}$ do not coincide and the dispersion curves are different. As is seen from the comparison with closed and unclosed strings, we deal with a topological effect.

It is usually considered that the boundaries must not affect the bulk properties of a system, since the volume of a near-boundary region is negligible relative to the total volume of a system. The smallness of the near-boundary region yields, in particular, the dimensional formulas of the form (3.1)–(3.3) from [38], in which the bulk characteristic (in [38], it is the ground-state energy per atom) of the system is invariable as $N, V \to \infty$, $N/V = const$. This implies that the shape of boundaries does not affect the bulk properties. We agree with this, it corresponds to the smallness of surface corrections A_j and Q_j in the above-obtained equations for Ψ_0 and $\Psi_{\mathbf{k}}$. Since any infinite system ($N, V = \infty$) has no boundaries, we make the second conclusion: the very presence of boundaries does not affect the bulk properties. One would think that this conclusion follows from the proofs [39] of the existence of a thermodynamic limit too. But we meet here the absence of a full rigor. A system with boundaries differs from an infinite system topologically: the latter has no boundaries and is topologically closer to a closed system. If the transition $N, V \to \infty$ means that to an infinite system, it is implicitly assumed that a change in the topology does not change the class of functions describing the bulk properties of a system. This should be proper for quantum systems at high T: the physical processes for them are conditioned by atoms, and an atom after the collision with a wall collides with other atoms and loses the memory of the wall. However, as $T \to 0$, the physics of systems is usually conditioned by waves (collective oscillations). The standing waves are modulated by walls and, therefore, "remember" the walls. Because the collections of harmonics for waves in an open and closed system are different, the bulk properties of these systems can be different as well. The above-presented calculation for He II confirms this. Since we deal with a topological effect, it should be valid not only for quantum fluids, but also for gases and crystals. In our opinion, the proofs of the existence of a thermodynamic limit mean only that it is possible to pass to the limit $N, V \to \infty$, $N/V = const$ in the formulas. But, in this case, we remain in the class of functions obtained for a finite system. If the infinite system is described by other functions, such a transition does not allow one to describe the infinite system, i.e., this transition is incorrect. Of course, such a problem does not arise for cyclic systems, because the transition $N, V \to \infty$ does not change their topology. The main point consists in that the effect is related to the topology of a system as a whole, rather than to the influence of a thin layer of near-surface atoms. In this meaning, the effect is a bulk one.

The role of BCs is also discussed in Appendix.

VI. THERMODYNAMICS

Under periodic BCs, the thermodynamic quantities are calculated from the free energy [40]

$$F = -k_B T \int \ln(1+n_q)\frac{d\mathbf{k}}{(2\pi)^3}. \qquad (75)$$

Under zero BCs, law (3) holds. Therefore, the quantity 2π in (75) should be replaced by π, which increases the value of the integral by eight times. However, if we change the sign of one or several components of $\mathbf{k}$ in the WF of a standing phonon, the WF is not changed. In other words, eight states with different signs of the components of $\mathbf{k}$ are equivalent. Therefore, we must integrate only over the sector $k_x, k_y, k_z > 0$, i.e., integral (75) decreases by eight times. We arrive at formula (75) and the known formulas for entropy, heat capacity, etc.

VII. POSSIBLE EXPERIMENTAL TESTS

Formula (73) implies that the dispersion curve of He II must be different for periodic and nonperiodic boundaries. In particular, it should be different for a monoatomic film of He II on a plane surface ($d = 2$ in Eq. (73)), on the lateral surface of a cylinder ($d = 1$), and on the surface of a torus ($d = 0$). Therefore, the heat capacity of these films at the same temperature must be significantly different. A more interesting effect is also possible. It is necessary to form a monoatomic film of He II on the surface of a torus and then to cut the torus (similarly to the breaking of a ring). As a result, some boundaries must arise. In this case, the system passes from the state with $d = 0$ to that with $d = 1$. Therefore, the energies of phonons and of the ground state are sharply changed, the ensemble of phonons will be rearranged, and the temperature of the system will jump by a value comparable with the initial T. If we restore the cut torus, the inverse processes will run. This is a phase transition that can be called a topological one. Its observation is quite real and can be a test for the effect of boundaries.

Our analysis shows that if the film thickness is more than several atomic layers, the properties of a helium film on the cylinder surface must be the same as those of helium in the cylinder bulk.

VIII. DISCUSSIONS

Of interest is the following property. The ground state is described by WF (26) with the product of N triple sines with the wave vector $\mathbf{k}_1$, and the excited state is presented by the same WF with the factor $\psi_{\mathbf{k}}$. The sines of the GS set a standing wave in the probability field. In the zero approximation, $\psi_{\mathbf{k}}$ is equal to $\rho_{-\mathbf{k}}$ with permutations and "is convolved" with the sines of the GS, by resulting in a superposition of states, whose structure coincides with that of Ψ_0 with the difference that the wave vector for some sines differs from $\mathbf{k}_1$. We may roughly say that *the excitation is simply the replacement of one of the standing waves of the GS by a wave with larger* $\boldsymbol{k}$. Such a structure of WFs means that the GS is formed by N identical standing "phonons" with the wave vector $\mathbf{k}_1$. In the excited states, a part of phonons is replaced by phonons with different $\mathbf{k}$. Therefore, we may say that the ground state and excited ones of the Bose fluid in a rectangular vessel contain N harmonics (phonons), and the difference consists in the composition of harmonics. In this case, the meaning of the notion of "excitation" is changed, because the GS becomes, in a certain sense, "excited", since it contains N harmonics with minimal $\mathbf{k}$.

With regard for corrections, the harmonics of the GS are not identical by structure to the harmonics of weakly excited states: the corrections for harmonics of the GS are collected in the exponent, whereas this is not the case for the excited states; and the equations for corrections are different. This is not amusing, because the GS is the resonant state with N identical interacting "phonons". Their structure must be different from the structure of single "ordinary" phonons described without regard for their interaction.

By passing through helium, neutrons create phonons. This process occurs with the conservation of the momentum. According to our solution, phonons are standing waves without definite momentum. However, the WF of a phonon is a sum of eight traveling waves, each with a definite momentum. Neutrons create, apparently, just such running waves. In a small region of helium, we see a lot of quasiparticles in the form of moving localized wave packets with a certain momentum. But if we consider helium on the whole, these quasiparticles turn out to be parts of the total system of standing waves filling the vessel.

We now return to the method. As was shown [17, 27], the collections $\rho_{\mathbf{k}_j}$ (19) are functionally independent, if $\mathbf{k}_j$ are quantized by the $2\pi/L$-law (2). But, in our sums, the wave vectors are quantized sometimes by the π/L-law (3); in this case, the collections $\rho_{\mathbf{k}_j}$ (19) are not functionally independent. For example, $\rho_{\mathbf{k}_j}$ with odd $\mathbf{k}$ (with components of the form $(2j+1)\pi/L$) can be expanded in collections (19) with even $\mathbf{k}$ (with components of the form $2j\pi/L$). But, in this case, we are faced with difficulties indicated below. Therefore, we find a solution without expansion. Such a means is quite correct. Indeed, let us seek a function in the form

$$F(x) = \sum_j A_j f_j(x), \tag{76}$$

and let the Schrödinger equation is reduced for it to the form

$$\sum_l B_l(\{A_j\}) f_l(x) = 0, \tag{77}$$

and we determine coefficients A_j such that $B_l = 0$ for all l. Then it is obvious that $F(x)$ is a solution irrespective of whether the functions $f_j(x)$ are independent and whether they form a full collection. We have acted just in this manner. As usual, the full collection is used in order that all solutions be found. But, under a certain structure of the equation, it is easier to find solutions if $f_j(x)$ have some particular structure and are dependent. It seems that this is our case. We have used the expansion in the variables $\rho_{\mathbf{k}_j}$ (19) with $\mathbf{k} \sim \pi j/L$ (3). Part of them, namely $\rho_{\mathbf{k}_j}$ (19) with even $\mathbf{k}$, forms the full collection, and the other variables are additional. Since the collection of functions, in which we expand, contains the full collection, no solutions should be missed.

We can try to use the independent basis functions as follows. Let us expand the potential in a series

$$\begin{aligned} \tfrac{1}{2}\sum_{i\neq j} U_3(|\mathbf{r}_i-\mathbf{r}_j|) &= c_1\varphi_1 + \sum_{\mathbf{k}\neq 0} c_2(\mathbf{k})\varphi_2 + \\ &+ \sum_{\mathbf{k}_1,\mathbf{k}_2\neq 0}^{\mathbf{k}_1+\mathbf{k}_2\neq 0} c_3(\mathbf{k}_1,\mathbf{k}_2)\varphi_3 + \ldots \end{aligned} \tag{78}$$

with $\mathbf{k}_j$ (2), where the functions [17]

$$\varphi_1 = A_1, \quad \varphi_2 = A_2(\rho_{\mathbf{k}}\rho_{-\mathbf{k}} - 1), \tag{79}$$

$$\begin{aligned} \varphi_3 &= \frac{A_3}{\sqrt{N}}\Big(\sqrt{N}\rho_{\mathbf{k}_1}\rho_{\mathbf{k}_2}\rho_{-\mathbf{k}_1-\mathbf{k}_2} + 2 - \\ &- (\rho_{\mathbf{k}_1}\rho_{-\mathbf{k}_1} + \rho_{\mathbf{k}_2}\rho_{-\mathbf{k}_2} + \rho_{\mathbf{k}_1+\mathbf{k}_2}\rho_{-\mathbf{k}_1-\mathbf{k}_2})\Big), \ldots \end{aligned} \tag{80}$$

($A_j = const$) form a complete orthonormalized basis for the expansion of any translation-invariant Bose symmetric function of the variables $\mathbf{r}_1, \ldots, \mathbf{r}_N$. However, such a method turns out not the best one: due to the presence of divergent terms, and because many basis functions include one-type terms of the form $\rho_{\mathbf{k}}\rho_{-\mathbf{k}}$, *etc.* such that it is difficult to gather and to evaluate them. In addition, the distinction of the zero and periodic BCs is insignificant in such an approach: for example, the coefficient c_2 differs only by a small relative value $\sim a/L$, where a is the interaction radius. For He II, we have $a/L \sim 1/N$. Therefore, in order to reveal the distinction of the BCs, we should consider $\sim N$ corrections in expansion (78), which is impossible to perform. In other words, the language of the functions φ_j is not suitable for taking boundaries into account.

Note also the following. The expansion in the $2\pi/L$-harmonics reproduces exactly the potential at the zero BCs at $|x_i - x_j|, |y_i - y_j|, |z_i - z_j| \leq L/2$. To specify the potential in the domain $> L/2$, we must additionally attract the odd π/L-harmonics from (15). In this case, the function is set by different collections of harmonics on different intervals. Therefore, the description with the use of only even $2\pi/L$-harmonics in the whole domain is not efficient and leads to the difficulties related to series (78). As opposite, the expansion (15) reproduces the potential in the entire domain on the basis of the single collection of harmonics and, therefore, is better.

IX. CONCLUSIONS

We have studied the influence of the walls of a vessel on the microstructure of He II and obtained a new class of exact solutions for the wave functions describing N interacting Bose particles in a box.

It is shown that the explicit account for boundaries changes the equations for the Bose liquid and the calculated values of the observed bulk quantities. Thus, the models with periodic boundary conditions should be reconsidered.

Probably, the wave function of the ground state of Bose crystal with rectangular lattice can be constructed [41] on the basis of the wave function (26) of Bose liquid.

Recently, we have solved the problem in an approach with independent basic functions [42]. In the limit of a weak interaction, the solution coincides with that obtained above. The further studies will help us to better understand how the boundaries affect the properties of Bose systems.

The author thanks N. Iorgov, K. Szalewicz and S. Yushchenko for valuable discussions.

X. APPENDIX A

It follows from (67) that the functions $Q_{l\geq 2}$ satisfy the symmetry relations

$$Q_2(\mathbf{q}, \mathbf{q}_1, \mathbf{k}) = Q_2(\mathbf{q}, -\mathbf{q}_1 - \mathbf{q} - \mathbf{k}, \mathbf{k}),$$

$$\begin{aligned} Q_3(\mathbf{q}, \mathbf{q}_1, \mathbf{q}_2, \mathbf{k}) &= Q_3(\mathbf{q}, \mathbf{q}_2, \mathbf{q}_1, \mathbf{k}) = \\ = \; & Q_3(\mathbf{q}, -\mathbf{q}_1 - \mathbf{q}_2 - \mathbf{q} - \mathbf{k}, \mathbf{q}_2, \mathbf{k}) = \\ = \; & Q_3(\mathbf{q}, \mathbf{q}_1, -\mathbf{q}_1 - \mathbf{q}_2 - \mathbf{q} - \mathbf{k}, \mathbf{k}), \end{aligned}$$

and the analogous relations can be written for higher Q_l.

Let us consider $b_0(\mathbf{k})$ (68). As seen from (68), $b_0(\mathbf{k})$ is nonzero, if all components of the vector $\mathbf{k}$ are multiple to $2\pi/L$. It follows from the equations that b_0 does not affect the dispersion curve $\epsilon(k)$. Since $\epsilon(k) \sim k k_{rot}$, relation (68) implies that, for the one-dimensional $\mathbf{k}$, we have $b_0 \sim N^{1/6}$ at $k \sim k_{rot}$. For the two- and three-dimensional $\mathbf{k}$, b_0 is less, respectively, by $N^{1/3}$ and $N^{2/3}$ times. For the irregular configurations of atoms, typical for a fluid, $\rho_{\mathbf{k}} \sim 1$. Therefore, for the one-dimensional $\mathbf{k}$, we have $b_0(\mathbf{k}) \gg \rho_{\mathbf{k}}$, i.e., b_0 determines the quantity $\psi^0_{\mathbf{k}}$ (67). At the normalization, b_0 "hides" itself, and we obtain $\psi_{\mathbf{k}} \approx 1$. By the structure of $\psi_{\mathbf{k}}$, such states are close to the ground one, but their energies are different at different k and larger than E_0. The nature of this anomaly is not clear. It can turn out fictitious, since $\Psi_{\mathbf{k}}$ (57), (60) with different $\mathbf{k}$ are not completely orthogonal due to the corrections with Q_l. These corrections cause a small mixing of different states: $\int d\Omega \Psi_{\mathbf{k}_1} \Psi^*_{\mathbf{k}_2 \neq \mathbf{k}_1} \sim N^{-1/3}$ at $\int d\Omega \Psi_{\mathbf{k}_1} \Psi^*_{\mathbf{k}_1} = 1$.

In addition to the regular solution, Eqs. (69)–(71) have a huge number of pole solutions related to the nullification of the denominators of the functions b_2, b_3, and Q_l in (69). For example, let we have a set of pole solutions of the form $\epsilon(k) = \epsilon_0(\mathbf{k}_2) + \epsilon_0(\mathbf{k} + \mathbf{k}_2) + \delta\epsilon(\mathbf{k}, \mathbf{k}_2)/N$ for a given k with different $\mathbf{k}_2$ and with $\delta\epsilon \sim 1$. It can be verified that these solutions disappear in the thermodynamic limit, but they form a new solution class for a system with finite size. However, these solutions are nonphysical. Indeed, we have $b_2(\mathbf{k}_2, \mathbf{k}) \sim N$ for pole $\mathbf{k}_2$. Hence, the term $\frac{b_2(\mathbf{k}_2,\mathbf{k})}{\sqrt{N}} \rho_{\mathbf{k}_2} \rho_{-\mathbf{k}_2-\mathbf{k}} \sim \sqrt{N} \rho_{\mathbf{k}_2} \rho_{-\mathbf{k}_2-\mathbf{k}}$, which is greater than the term $\rho_{-\mathbf{k}}$ by $\sim \sqrt{N}$ times, becomes the main term ($\sim \sqrt{N}$) in expansion (58). The quantity $\psi_{\mathbf{k}}$ corresponds to distorted two-phonon state. The same is true for poles of Q_2. The poles of b_3 correspond to distorted three-phonon states. Moreover, the poles of Q_1 correspond to a phonon with a different wave vector.

XI. APPENDIX B. MODEL OF POINT PARTICLES

The explicit account for boundaries was performed within the exactly solvable one-dimensional model [12, 13] for bosons with the point interaction (point particles with the interatomic potential as the δ-function). For such a system, the periodic BCs turn out, in essence, to be equivalent to the zero BCs: the values of E_0 for both BCs differ from each other only by a negligible value $\sim E_0/N$. At the same time, it was obtained above for the Bose fluid that the values of E_0 differ strongly under periodic and zero BCs.

Such a difference may be caused by the difference of interatomic potentials. In our opinion, the principal point consists in the following. Consider a cyclic system. The solution for the WFs of point particles is [38]:

$$\Psi_{(k)}(x) = \sum_P a(P) P e^{i \sum\limits_{l=1}^{N} k_{P_l} x_l}, \tag{81}$$

where P are all permutations of k_l. WF (81) is a superposition of states of N free particles. The energy,

$$E = (\hbar^2/2m)(k_1^2 + k_2^2 + \ldots + k_N^2), \tag{82}$$

is the same for all these states. At the nonpoint interaction, the WF of the ground state is given by (4), (5).

If we expand this WF in a Fourier series, we obtain a superposition of states (in 1D)

$$\Psi_0(x) = \sum_P a(P(\{x_l\})) \sum_{k_1\ldots k_N} A(k_1,\ldots,k_N) e^{i\sum\limits_{l=1}^{N} k_l x_l}, \tag{83}$$

with $k_l = 2\pi l/L$ and energy (82), which is different for different terms. This Ψ_0 cannot be presented in the form of a superposition of plane waves with the same energy.

In the presence of walls, a WF of point particles was constructed [12] by the optical analogy as a sum of plane waves (81) and their all reflections with the same energy. Therefore, a wall does not change the energy of the system. According to [12], the solutions for N particles in a vessel with length L coincide with definite solutions for $2N$ particles in a periodic vessel $2L$ in length. However, the waves in (83) have different energies, and the addition of reflected waves will lead to the redistribution of terms and to a change in the energy of the system. This can be seen at another angle of vision, if we take into account that, according to (4) and (5), $\Psi_0 = \exp(\sum\limits_{\mathbf{k}} a_2(\mathbf{k})\rho_{\mathbf{k}}\rho_{-\mathbf{k}}/2)$ is the WF of the ground state of oscillators with coordinates $\rho_{\mathbf{k}}$, and the corrections $a_{j\geq 3}$ are related to the interaction of oscillators. Let us have a pair of oscillators with the same frequency ω_1. If we add to them a pair of the same oscillators, which does not interact with the first pair, then the frequency of all oscillators remains to be ω_1. This corresponds to the solution for a point interaction. But if we include the interaction between pairs, then the eigenfrequencies of the system are changed, as is known from mechanics. Instead of two frequencies, we will have four ones, or more. This case corresponds to a nonpoint interaction.

The absence of the effect of boundaries under a point interaction can be imagine, probably, in the following way: The effect is related to a certain game of harmonics; however, the point interaction has no dispersion ($\nu(k) = const$), and no game is present.

We have tried to qualitatively explain the effect. But these analogies are rather rough, and only the exact solution of the problem can be a proof.

Since Ψ_0 of the systems with point and nonpoint interactions are different, the solutions for E_0 and $E(k)$ should be different too. However, it turns out that the solutions [12, 13, 38] for a weak point interaction are close to Bogolyubov ones. This is related to the fact that Bogolyubov approach does not use WFs, and it is possible to describe the systems with point and nonpoint interactions in the same way (in the first case, the potential can be considered in the $\mathbf{r}$-space not passing to Fourier transforms). Therefore, it is of importance to find the solutions for a noncyclic nonpoint system namely in Bogolyubov approach and to determine to which solutions they correspond: to the traditional solutions for a cyclic system or to the above-obtained ones. We now study this problem.

[1] L. Landau, J. Phys. USSR **5**, 71 (1941); **11**, 91 (1947).
[2] N.N. Bogoliubov, J. Phys. USSR **11**, 23 (1947).
[3] N.N. Bogoliubov and D.N. Zubarev, Sov. Phys. JETP **1**, 83 (1956).
[4] R. Feynman, Phys. Rev. **94**, 262 (1954).
[5] R.P. Feynman and M. Cohen, Phys. Rev. **102**, 1189 (1956).
[6] H.R. Glyde and E.C. Svensson, in *Methods of Experimental Physics*, vol. 23, part B, edited by D.L. Price and K. Skold (Academic Press, New York, 1987), p. 303.
[7] L. Reatto, J. Low Temp. Phys. **87**, 375 (1992).
[8] M.D. Tomchenko, arXiv:cond-mat/0904.4434.
[9] A. Bijl, Physica **7**, 869 (1940).
[10] M.D. Girardeau, J. Math. Phys. **6**, 1083 (1965).
[11] K. Huang, *Statistical Mechanics* (Wiley, New York, 1987), Chapter 19.
[12] M. Gaudin, Phys. Rev. A **4**, 386 (1971).
[13] M. Gaudin, *La Fonction d'Onde de Bethe* (Masson, Paris, 1983).
[14] M. Tomchenko, arXiv:cond-mat/1201.2185.
[15] C.E. Campbell, Phys. Lett. **44A**, 471 (1973).
[16] E. Feenberg, Ann. Phys. **84**, 128 (1974).
[17] I.A. Vakarchuk and I.R. Yukhnovskii, Theor. Math. Phys. **40**, 626 (1979).
[18] P.A. Whitlock, D.M. Ceperley, G.V. Chester, and M.H. Kalos, Phys. Rev. B **19**, 5598 (1979).
[19] M.H. Kalos, M.A. Lee, P.A. Whitlock, and G.V. Chester, Phys. Rev. B **24**, 115 (1981).
[20] E. Krotscheck, Phys. Rev. B **33**, 3158 (1986).
[21] J.L. Epstein and E. Krotscheck, Phys. Rev. B **37**, 1666 (1988).
[22] T. MacFarland, S.A. Vitiello, L. Reatto, G.V. Chester, and M.H. Kalos, Phys. Rev. B **50**, 13577 (1994).
[23] L. Reatto, G.L. Masserini, and S.A. Vitiello, Physica B **197**, 189 (1994).
[24] I.O. Vakarchuk, V.V. Babin, and A.A. Rovenchak, J. Phys. Stud. (Lviv) **4**, 16 (2000).
[25] M. Tomchenko, Low Temp. Phys. **32**, 38 (2006).
[26] R. Rota, J. Boronat, J. Low Temp. Phys. **166**, 21 (2012).
[27] I.A. Vakarchuk and I.R. Yukhnovskii, Theor. Math. Phys. **42**, 73 (1980).
[28] M.D. Tomchenko, Ukr. J. Phys. **50**, 720 (2005).
[29] E. Zaremba and W. Kohn, Phys. Rev. B **15**, 1769 (1977).
[30] K.V. Grigorishin, B.I. Lev, Ukr. J. Phys. **56**, 1182 (2011).
[31] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics. Non-Relativistic Theory* (Pergamon Press, New York, 1980).
[32] S.A. Yushchenko, private communication.
[33] A.R. Jansen and R.A. Aziz, J. Chem. Phys. **107**, 914 (1997).
[34] M. Przybytek, W. Cencek, J. Komasa, G. Lach, B. Jeziorski, and K. Szalewicz, Phys. Rev. Lett. **104**, 183003 (2010); **108**, 129902 (E) (2012).
[35] M.R. Andrews, D.M. Kurn, H.-J. Miesner, D.S. Durfee,

C.G. Townsend, S. Inouye, and W. Ketterle, Phys. Rev. Lett. **79**, 553 (1997).

[36] L.P. Pitaevskii, Sov. Phys. JETP **9**, 830 (1959).

[37] J.M. Ziman, *Principles of the Theory of Solids* (Cambridge Univ. Press, Cambridge, 1964), Chapter 1.

[38] E.H. Lieb, W. Liniger, Phys. Rev. **130**, 1605 (1963); E.H. Lieb, Phys. Rev. **130**, 1616 (1963).

[39] D. Ruelle, *Statistical Mechanics. Rigorous Results* (Benjamin Inc., New York, 1969).

[40] I.M. Khalatnikov, *An Introduction to the Theory of Superfluidity* (Perseus, Cambridge, 2000), Chapter 1.

[41] M. Tomchenko, arXiv:cond-mat/1201.2623.

[42] M. Tomchenko, arXiv:cond-mat/1204.2149.